

09/220,398

FILE 'HOME' ENTERED AT 07:46:28 ON 02 DEC 2001

=> file biosis medline caplus wpids
COST IN U.S. DOLLARS

SINCE FILE ENTRY	TOTAL SESSION
0.15	0.15

FULL ESTIMATED COST

FILE 'BIOSIS' ENTERED AT 07:46:50 ON 02 DEC 2001
COPYRIGHT (C) 2001 BIOSIS(R)

FILE 'MEDLINE' ENTERED AT 07:46:50 ON 02 DEC 2001

FILE 'CAPLUS' ENTERED AT 07:46:50 ON 02 DEC 2001
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'WPIDS' ENTERED AT 07:46:50 ON 02 DEC 2001
COPYRIGHT (C) 2001 DERWENT INFORMATION LTD

*** YOU HAVE NEW MAIL ***

=> s linking nucleic acid
L1 11 LINKING NUCLEIC ACID

=> s l1 and multicomponent
L2 2 L1 AND MULTICOMPONENT

=> d l2 bib abs

L2 ANSWER 1 OF 2 BIOSIS COPYRIGHT 2001 BIOSIS
AN 2001:473230 BIOSIS
DN PREV200100473230
TI Method and kits for preparing **multicomponent** nucleic acid
constructs.
AU Harney, Peter D. (1)
CS (1) Aliso Viejo, CA USA
ASSIGNEE: VectorObjects, LLC, Wellesley, MA, USA
PI US 6277632 August 21, 2001
SO Official Gazette of the United States Patent and Trademark Office Patents,
(Aug. 21, 2001) Vol. 1249, No. 3, pp. No Pagination. e-file.
ISSN: 0098-1133.
DT Patent
LA English
AB The invention provides a highly efficient, rapid, and cost effective
method of **linking nucleic acid** components in
a predetermined order to produce a nucleic acid **multicomponent**
construct. The invention further provides nucleic acid components, each
nucleic acid component comprising a double stranded nucleic acid molecule
having at least one single stranded 5' or 3' terminal sequence, the
terminal sequence having sufficient complementarity to either a terminal
sequence in a separate nucleic acid component or to a sequence in a
linking nucleic acid molecule so as to allow
for specific annealing of complementary sequences and linkage of the
components in a predetermined order. Kits containing reagents required to
practice the method of the invention are also provided.

=> d l2 bib abs 2

L2 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2001 ACS
AN 1998:31322 CAPLUS
DN 128:85140

TI Preparation of **multicomponent** nucleic acid constructs by
 ligation of nucleic acid components with complementary terminal sequences
 IN Harney, Peter D.
 PA Biodynamics Associates, USA
 SO PCT Int. Appl., 33 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9748716	A1	19971224	WO 1997-US10523	19970616
	W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
	CA 2258570	AA	19971224	CA 1997-2258570	19970616
	AU 9733997	A1	19980107	AU 1997-33997	19970616
	EP 915903	A1	19990519	EP 1997-930083	19970616
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
	JP 2000512852	T2	20001003	JP 1998-503272	19970616
	US 6277632	B1	20010821	US 1997-877034	19970616
PRAI	US 1996-19869	P	19960617		
	WO 1997-US10523	W	19970616		
AB	<p> The invention provides a highly efficient, rapid, and cost effective method of linking nucleic acid components in a predetd. order to produce a nucleic acid multicomponent construct. The invention further provides nucleic acid components, each nucleic acid component comprising a double-stranded nucleic acid mol. having at least one single-stranded 5' or 3' terminal sequence, the terminal sequence having sufficient complementarity to either a terminal sequence in a sep. nucleic acid component or a sequence in a linking nucleic acid mol. so as to allow for specific annealing of complementary sequences and linkage of the components in a predetd. order. The various nucleic acid components can be linked via, without limitation, the following: (1) annealing of 5' complementary terminal sequences in 2 sep. nucleic acid components; (2) annealing of 3' complementary terminal sequences in 2 sep. nucleic acid components; (3) annealing of an oligonucleotide bridge with complementary 5' and 3' terminal sequences in 2 sep. nucleic acid components; or (4) annealing of an adaptor mol. with complementary 5' or 3' terminal sequences in 2 sep. nucleic acid components;. To demonstrate the simultaneous assembly of multiple nucleic acid components having unique, non-palindromic terminal sequences, to produce a viable plasmid vector, 3 nucleic acid components were used: a gene coding for green fluorescent protein; a 0.6-kb mol. coding for terminator sequences and a histidine tag; and a 2.5-kb mol. coding for the lac promoter, an ampicillin resistance gene, and an origin of replication. This method is particularly suitable for the construction of nucleic acid vectors. </p>				

=>